

Gulf of Mexico Harmful Algal Bloom Bulletin

Region: Texas


Monday, 19 December 2016

NOAA National Ocean Service

NOAA Satellite and Information Service

NOAA National Weather Service

Last bulletin: Monday, December 12, 2016



Satellite chlorophyll image with possible *K. brevis* HAB areas shown by red polygon(s), when applicable. Points represent cell concentration sampling data from December 9 to 15: red (high), orange (medium), yellow (low b), brown (low a), blue (very low b), purple (very low a), pink (present), and green (not present). Cell count data are provided by Texas Parks and Wildlife Department. For a list of sample providers and a key to the cell concentration categories, please see the HAB-OFS bulletin guide:

http://tidesandcurrents.noaa.gov/hab/hab_publication/habfs_bulletin_guide.pdf

Detailed sample information can be obtained through the Texas Parks and Wildlife Department at: http://www.tpwd.state.tx.us./landwater/water/environconcerns/hab/redtide/status.phtml

Conditions Report

Karenia brevis (commonly known as Texas red tide) ranges from not present to very low concentrations along the coast of Texas. No respiratory irritation is expected alongshore Texas Monday, December 19 through Tuesday, December 27.

Check http://tidesandcurrents.noaa.gov/hab/beach_conditions.html for recent, local observations.

Analysis

Due to the upcoming federal holiday, the next bulletin will be issued on Tuesday, December 27.


Sampling from the Texas A&M University's Imaging FlowCytobot, located on the Port Aransas ship channel, indicates that *Karenia brevis* concentrations range between 'not present' and 'very low a' (TAMU; 12/12-19). For information on area shellfish restrictions, contact the Texas Department of State Health Services.

Recent ensemble imagery (MODIS Aqua, 12/16; shown left) is obscured by clouds north of the Padre Island National Seashore (PINS) region, limiting analysis. Elevated chlorophyll (2-6 μ g/L) is visible along- and offshore from PINS to south of the Rio Grande. Elevated chlorophyll is not indicative of the presence of *K. brevis* and is most likely due to the resuspension of benthic chlorophyll and sediments along the coast.

Forecast models based on predicted near-surface currents indicate a potential maximum transport of greater than 150 km south from the Port Aransas region from December 16-22.

Kavanaugh, Urízar

To see previous bulletins and forecasts for other Harmful Algal Bloom Bulletin regions, visit the NOAA Harmful Algal Bloom Operational Forecast System bulletin archive: http://tidesandcurrents.noaa.gov/hab/bulletins.html




Wind speed and direction are averaged over 12 hours from buoy measurements. Length of line indicates speed; angle indicates direction. Red indicates that the wind direction favors upwelling near the coast. Values to the left of the dotted vertical line are measured values; values to the right are forecasts. Wind observation and forecast data provided by NOAA's National Weather Service (NWS).

-2-

Wind Analysis

Port Aransas to Matagorda Ship Channel: North winds (5-25kn, 3-13m/s) today through Tuesday night. Northeast to east winds (5kn, 3m/s) Wednesday through Wednesday night. North winds (15-25kn, 8-10m/s) after midnight Wednesday night through Thursday night. Northeast to east winds (10-15kn, 5-8m/s) after midnight Thursday night through Friday night. Southeast winds after midnight Friday night.



Verified and suspected HAB areas shown in red. Other areas with *K. brevis* optical characteristics shown in yellow (see p. 1 analysis for interpretation).